

# Interspecific and intraspecific transference of metabolism-based mesotrione resistance in dioecious weedy *Amaranthus*

Maxwel C. Oliveira<sup>1,\*</sup> , Todd A. Gaines<sup>2</sup> , Eric L. Patterson<sup>2</sup>, Amit J. Jhala<sup>1</sup>, Suat Irmak<sup>3</sup>, Keenan Amundsen<sup>1</sup> and Stevan Z. Knezevic<sup>1</sup>

<sup>1</sup>Department of Agronomy and Horticulture, University of Nebraska-Lincoln, Lincoln, NE, USA,

<sup>2</sup>Department of Bioagricultural Sciences and Pest Management, Colorado State University, Fort Collins, CO, USA, and

<sup>3</sup>Department of Biological Systems Engineering, University of Nebraska-Lincoln, Lincoln, NE, USA

Received 16 May 2018; revised 1 August 2018; accepted 6 September 2018; published online 11 September 2018.

\*For correspondence (e-mail max.oliveira@wisc.edu).

## SUMMARY

Pollen-mediated gene flow (PMGF) might play an important role in dispersing herbicide resistance alleles in dioecious weedy *Amaranthus* species. Field experiments in a concentric donor–receptor design were conducted to quantify two sets of PMGF studies, an interspecific (*Amaranthus tuberculatus* × *Amaranthus palmeri*) and an intraspecific (*A. tuberculatus* × *A. tuberculatus*). In both studies, PMGF was evaluated using a resistant *A. tuberculatus* phenotype with enhanced mesotrione detoxification via P450 enzymes as a source of resistance alleles. For interspecific hybridization, more than 104 000 putative hybrid seedlings were screened with three markers, one phenotypic and two molecular. The two molecular markers used, including 2-bp polymorphisms in the internal transcribed spacer region, distinguished *A. palmeri*, *A. tuberculatus* and their hybrids. Results showed that 0.1% hybridization between *A. tuberculatus* × *A. palmeri* occurred under field research conditions. For intraspecific hybridization, 22 582 seedlings were screened to assess the frequency of gene flow. The frequency of gene flow ( $F_{GF}$ ) varied with distance, direction and year of the study. The farthest distance for 90% reduction of  $F_{GF}$  was at 69 m in 2015 however, after averaging across directions it was 13.1 and 26.1 m in 2014 and 2015, respectively. This study highlights the transfer of metabolism-based mesotrione resistance from *A. tuberculatus* to *A. palmeri* under field research conditions. The results presented here might aid in the rapid detection of *A. palmeri* among other *Amaranthus* species and show that PMFG could be expediting the increase of herbicide resistance in *A. palmeri* and *A. tuberculatus* across US crop production areas.

**Keywords:** *Amaranthus palmeri*, *Amaranthus tuberculatus*, gene flow, herbicide resistance evolution, HPPD-inhibitor herbicide, hybridization, Palmer amaranth, waterhemp.

## INTRODUCTION

The dioecious weedy *Amaranthus* species *Amaranthus palmeri* and *Amaranthus tuberculatus* are currently the most economically important weed species infesting row crop areas in the southern and north-central USA, respectively (Steckel, 2007; Webster and Nichols, 2012; Ward *et al.*, 2013). Various farm practices have contributed to the adaptation of *Amaranthus* species to modern cropping systems, particularly the reduction in the use of soil-applied herbicide and increased adoption of no-tillage row crop production systems (Culpepper, 2006; Owen, 2008). Additionally, these species have biological characteristics that increase their ability to be problem weeds, including dioecism and high fecundity (Hartzler *et al.*, 2004; Refsell and Hartzler,

2009; Ward *et al.*, 2013). The *Amaranthus* genus is notorious for a high rate of spontaneous intraspecific and interspecific hybridization (Grant, 1959; Greizerstein and Poggio, 1995; Ellstrand *et al.*, 1996). Dioecism in *Amaranthus* species is an important component of adaptation and evolution of these species, and may partially explain the high level of herbicide resistance existing across this genus. Populations of *A. palmeri* and *A. tuberculatus* have evolved resistance to six herbicide sites of action (SOA) throughout the USA (Tranel *et al.*, 2011; Bernards *et al.*, 2012; Ward *et al.*, 2013). The high frequency of herbicide resistance in this genus, dioecy, large population sizes and intense selection pressure make *Amaranthus* an excellent model for studying plant evolution at the landscape level.

Herbicide resistance alleles might be spread via pollen-mediated gene flow (PMGF) (Jasieniuk *et al.*, 1996). Liu *et al.* (2012) reported that pollen from an *A. tuberculatus* population has great longevity and long-distance dispersal. As a result, there is the potential for intraspecific (Sosnoskie *et al.*, 2012; Sarangi *et al.*, 2017) and interspecific hybridization in *Amaranthus* species (Tranel *et al.*, 2002; Trucco *et al.*, 2005, 2006), transferring herbicide resistance traits (Gaines *et al.*, 2012). Therefore, PMGF might be an important aspect of the transfer of herbicide resistance genes, and it is likely to increase the adaptation success of dioecious *Amaranthus* species in modern cropping-systems.

Several molecular markers have been developed to distinguish *Amaranthus* species and their interspecific hybrids (Wetzell *et al.*, 1999a; Tranel *et al.*, 2002; Wright *et al.*, 2016); however, most markers rely on restriction fragment length polymorphisms (RFLPs), which can be expensive and time-consuming when genotyping large number of plants. More recently, a quantitative marker has been developed to identify *A. palmeri* from other *Amaranthus* species in mixed seed collections (Murphy *et al.*, 2017). This assay is particularly useful as *A. palmeri* has recently been listed as a noxious weed in several states in the north-central USA. Molecular assays for rapid identification of *Amaranthus* species and their hybrids are necessary for the identification and management of *Amaranthus* species.

While molecular markers exist for determining certain hybrids, most intraspecific PMGF studies in weedy *Amaranthus* are based on phenotypic herbicide resistance traits (Liu *et al.*, 2017; Sarangi *et al.*, 2017). Target-site resistance (TSR) mechanisms are usually conferred by a single dominant gene and can be used as a robust marker for identifying intraspecific hybridization (Mallory-Smith *et al.*, 2015). However, non-target-site resistance (NTSR) mechanisms have also recently been reported in *A. tuberculatus* and *A. palmeri* populations (Evans *et al.*, 2017; Figueiredo *et al.*, 2017; Kaundun *et al.*, 2017; Nakka *et al.*, 2017; Küpper *et al.*, 2018); some of these have complex inheritance (Hausman *et al.*, 2011) or the resistance gene has not yet been determined (Délye, 2013a). Recently there has been an increased interest in deciphering the genetic basis of NTSR mechanisms (Délye *et al.*, 2013b), but our knowledge of this area is far from complete (Gaines *et al.*, 2014; Duhoux *et al.*, 2015; Pan *et al.*, 2016). Even though PMGF is one of the key components for understanding ecological and evolutionary dynamics of NSTR mechanisms, the role of expression of NTSR herbicide resistance alleles in interspecific and intraspecific hybrids or  $F_1$  individuals under field research conditions remains to be explored.

Morphological similarities and potential hybridization between *Amaranthus* species are likely to increase the complexity of identification and management (Wassom and Tranel, 2005); therefore, it is critical to develop

molecular markers that robustly, effectively and accurately detect *Amaranthus* species. We have developed two competitive allele-specific PCR (KASP) markers (He *et al.*, 2014) to distinguish *A. palmeri*, *A. tuberculatus* and their interspecific hybrid. We hypothesized that: (i) PMGF with herbicide resistance transfer would occur from mesotrione-resistant (R) *A. tuberculatus* to mesotrione-susceptible (S) *A. palmeri* populations (interspecific hybridization) and (ii) PMGF transferring herbicide resistance would occur from R *A. tuberculatus* to S *A. tuberculatus* populations (intraspecific hybridization) under field conditions.

## RESULTS

### Interspecific (R *A. tuberculatus* × S *A. palmeri*) hybridization

More than 104 000 putative hybrid (R *A. tuberculatus* × S *A. palmeri*) seedlings were screened with 185 g active ingredient (ai) ha<sup>-1</sup> (1.75 times the mesotrione field rate), which was used as a cutoff rate expected to keep false positives low. Only 2.4% and 0.9% of putative hybrid seedlings survived mesotrione application in 2014 and 2015, respectively (Table 1).

Two KASP molecular marker assays were employed to test putative hybrids and eliminate false positive hybrids that survived the mesotrione screening test. The first marker was a single nucleotide polymorphism (SNP) in the acetolactate synthase (ALS) gene (ALS-SNP) and the second marker was a 2-bp polymorphism in the internal transcribed spacer (ITS) of the ribosomal coding region (TBP-ITS). The R *A. tuberculatus* and S *A. palmeri* parents clustered at their respective genotypes, except for a single S *A. palmeri* individual for the ALS-SNP that clustered in the no template control (NTC; Figure 1). The majority (79 out 86) of putative hybrids that survived mesotrione application clustered in the S *A. palmeri* region with only seven hybrids classified as heterozygotes for both SNPs (Figure 1). The rate of hybrids using KASP molecular assays was 0.042 and 0.128 for 2014 and 2015, respectively. These numbers were further used to calculate the frequency of interspecific hybridization.

In 2014, interspecific hybrids were found in the north-east (0.5 m) and north-west (5 m) block. In 2015, hybrids were found south (0.5 m), two east (0.5 m) and two at 15 m (north-east and north-west) of the R *A. tuberculatus* pollen-source block. The total estimated hybridization frequency was similar for both years (0.001), and power analysis confirmed the experiment's precision for detecting hybrids (Table 1).

### Intraspecific (R *A. tuberculatus* × S *A. tuberculatus*) hybridization

The model predicted the frequency of gene flow ( $F_{GF}$ ;  $F_1$  individuals) to be highest near the pollen-source block

**Table 1** Frequency of interspecific pollen-mediated gene flow (PMGF) between resistant *Amaranthus tuberculatus* and susceptible *Amaranthus palmeri* at different distances (pooled directions) under field research conditions at the Haskell Agricultural Laboratory of the University of Nebraska–Lincoln in 2014 and 2015.

Year	Distance <sup>a</sup>	Mesotrione treatment <sup>b</sup>		KASP assays <sup>c</sup>		Hybridization <sup>d</sup>	
		Emerged plants (E)	Living plants (A)	Plants analyzed	Hybrids	Estimated	Frequency
	m	#					
2014	0	4154	215	4	0	9	0.0022
	0.5	15 546	471	25	0	20	0.0013
	2	11 879	351	8	1	15	0.0013
	5	13 083	394	3	1	17	0.0013
	15	12 696	162	1	0	7	0.0005
	30	8351	86	3	0	4	0.0004
	45	8822	89	3	0	4	0.0004
	Total	74 531	1768	47	2	75	0.0010
2015	0	1044	25	1	0	3	0.0031
	0.5	5998	107	11	3	14	0.0023
	2	6050	47	5	0	6	0.0010
	5	6449	57	6	0	7	0.0011
	15	5297	49	6	2	6	0.0012
	30	8351	40	5	0	5	0.0006
	45	4889	13	5	0	2	0.0003
	Total	38 078	338	39	5	43	0.0011

<sup>a</sup>Distance of mesotrione-susceptible *A. palmeri* female plants from the pollen-source block of mesotrione-resistant *A. tuberculatus*.

<sup>b</sup>Number of emerged (E) and living (A) plants (putative hybrids, resistant *A. tuberculatus* × susceptible *A. palmeri*) treated with 185 g ai ha<sup>-1</sup> (1.75 times the mesotrione field rate) in interspecific hybridization only.

<sup>c</sup>Subsamples of living plants (A) from mesotrione treatment. The frequency of hybridization detected in the KASP assays (T) was 0.042 (2/47) and 0.128 (5/39) for 2014 and 2015, respectively. Hybrids were detected using two molecular KASP assays.

<sup>d</sup>The frequency of hybridization was calculated by combining phenotypic and molecular markers as described in equation (3) (frequency of hybridization = TA/E). Estimation of the number of hybrids was calculated from frequency of hybridization × living plants (A). A power analysis using binomial probabilities was 0.99 of the theoretical frequency of 0.5% hybridization at  $\alpha = 0.05$ .

(0 m) and decrease exponentially with distance (Figure 2). The maximum  $F_{GF}$  at 0.5 m was 0.67 in 2015, which is below the expected random mating value (1) and just over twice the  $F_{GF}$  in 2014 (0.32) (Table 2). The modeling efficiency (ME) of predicted  $F_{GF}$  ranged from 0.12 to 0.71, and the ME averaged lower than 0.40.

Female *S. A. tuberculatus* plants were not included inside the pollen-source block; therefore,  $F_{GF}$  at 0 m was not determined. However, the double-exponential model predicted  $F_{GF}$  at 0 m for each year and cardinal direction (Figure S1 in the online Supporting Information). The  $F_{GF}$  at 0 m aided in predicting distance at 50% and 90%  $F_{GF}$  reduction (Figure 3). The  $F_{GF}$  reduction values varied according to direction. In 2014, the distance for 90%  $F_{GF}$  reduction ranged from 1.5 to 19 m; however, in 2015, values varied from 4 to 69 m.


The average wind speed during the 6-week pollination period was lower than 6 m s<sup>-1</sup> blowing from all directions, but with higher wind frequencies from the south in both years (Figure 4). Wind frequency did not correlate with  $F_{GF}$ , but the frequency of wind direction was variable across

weeks in 2014 and 2015 (Figures S2 and S3). Therefore, the wind was blowing from different directions in different weeks, and typically varied within a day. The *Amaranthus* species in the present study are dioecious and flowering time varied within 6 weeks from the earliest to latest blooming plants, and even plants in the same distance/direction showed different flowering patterns.

## DISCUSSION

### Interspecific (R *A. tuberculatus* × S *A. palmeri*) hybridization

*Amaranthus palmeri* is described as the most economically damaging weed of southern US cropping-systems (Ward *et al.*, 2013). In recent years, *A. palmeri* has migrated into north-central states, overlapping in territory with *A. tuberculatus*, a major weed existing in north-central cropping systems (Kohrt *et al.*, 2017). Here, we failed to reject our hypotheses, demonstrating that transference of NTSR mechanisms of mesotrione resistance from R *A. tuberculatus* to S *A. palmeri* occurs under field research conditions.




**Figure 1.** Cluster analysis of competitive allele-specific PCR (KASP) assays of suspected hybrids (*A. tuberculatus* × *A. palmeri*) with two molecular markers. An acetolactate synthase single nucleotide polymorphism (ALS-SNP) marker (left) and a 2-bp polymorphism in the internal transcribed spacer (TBP-ITS) marker (right). Parental *Amaranthus tuberculatus* and *Amaranthus palmeri* were used as a positive control and no template for the negative control (NTC). Dashed lines represent cutoffs for making genotyping calls. The solid quarter circle is the cutoff for no amplification. Cutoffs were determined based on the *K*-means cluster analysis. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)].

Therefore, the complexity of managing weeds of these dioecious *Amaranthus* species is likely to increase in row crops of the north-central USA due to species coexistence in this region and potential hybridization between *A. tuberculatus* and *A. palmeri*.

The fitness penalties of R *A. tuberculatus* and S *A. palmeri* hybrids were not the objective in our study. Nonetheless, transfer of *A. palmeri* herbicide resistance genes to *A. tuberculatus* has been previously documented, showing apparent normal hybrid growth and fertility (Wetzel *et al.*, 1999b; Franssen *et al.*, 2001a,b). However, other studies have demonstrated negative effects in the *A. tuberculatus* and *A. palmeri* (reciprocal cross) hybrids, including irregular crossing over in meiosis (Steinau *et al.*, 2003) and triploid apomictic seed (Trucco *et al.*, 2007). The majority of weedy *Amaranthus* species are diploid ( $2n$ ) with their chromosome number ranging from 32 to 34 (Trucco *et al.*, 2007; Gaines *et al.*, 2012). Similar chromosome numbers could facilitate hybridization in *Amaranthus* species (Trucco *et al.*, 2005; Nandula *et al.*, 2014). While *A. tuberculatus* is shown to have 32 chromosomes, *A. palmeri* was documented with 34 chromosomes (Grant, 1959; Gaines *et al.*, 2012). In our study, R *A. tuberculatus* and S *A. palmeri* hybrid formation was low (0.001), which might be due to the differential chromosome number between these two species. Additional research to investigate the potential fitness trade-offs associated with *A. palmeri* and *A. tuberculatus* hybrids carrying NTSR mechanisms is necessary, including the possible balance of benefits of

herbicide resistance from gene flow with fitness penalties due to hybridization.

The expansion of *A. palmeri* into the north-central USA means it is currently sharing the same habitat with several other weedy *Amaranthus* species, including *Amaranthus spinosus* ( $2n = 34$ ), *Amaranthus powellii* ( $2n = 34$ ), *Amaranthus retroflexus* ( $2n = 34$ ) and *Amaranthus hybridus* ( $2n = 32$ ) (Sauer, 1957, 1967; Grant, 1959; Wetzel *et al.*, 1999a). Additionally, 10 *Amaranthus* species are dioecious, which aids cross-pollination (Steckel, 2007). Pollen in *Amaranthus* can spread for long distances by wind dispersal, which can increase the chances of hybridization among species and spread herbicide resistance alleles (Franssen *et al.*, 2001b; Sosnoskie *et al.*, 2009; Liu *et al.*, 2012). In some cases, hybridization in *Amaranthus* species might be unidirectional. For example, Trucco *et al.* (2009) showed that *A. hybridus* can transfer herbicide resistance alleles to *A. tuberculatus* but the reciprocal genetic exchange was not possible. Nonetheless, transfer of glyphosate resistance genes was also documented from *A. palmeri* to *A. tuberculatus*, *A. hybridus* and *A. spinosus* (Gaines *et al.*, 2012; Nandula *et al.*, 2014). Many of the *Amaranthus* species have similar morphologies and coexist in the same location, and their ability to hybridize makes it difficult to clearly differentiate them on the basis of morphology. As we have demonstrated, robust molecular KASP assays (ALS-SNP and TBP-ITS) can be used to distinguish R *A. tuberculatus*, S *A. palmeri* and their hybrids.



**Figure 2.** Intraspecific frequency of gene flow. Intraspecific frequency of gene flow (mesotrione-resistant *Amaranthus tuberculatus* × mesotrione-susceptible *A. tuberculatus*) affected by distance and eight cardinal directions (N, NE, E, SE, S, SW, W, NW) in a field research experiment at the Haskell Laboratory of the University of Nebraska–Lincoln. Modeling efficiency (ME) for 2014 and 2015. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)].

**Table 2** Frequency of intraspecific pollen-mediated gene flow (PMGF) in  $F_1$  individuals (resistant *Amaranthus tuberculatus* × susceptible *A. tuberculatus*) under field research conditions at Haskell Agricultural Laboratory of the University of Nebraska-Lincoln in 2014 and 2015

Year	Distance <sup>a</sup> m	Mesotrione treatment <sup>b</sup>		$F_1$ individuals <sup>c</sup>	
		Emerged plants #	Living plants	Frequency	Power <sup>d</sup>
2014	0.5	574	185	0.32	0.96
	2	605	135	0.22	1.00
	5	725	103	0.14	1.00
	15	570	30	0.05	0.99
	30	611	24	0.04	0.97
	45	572	11	0.02	0.40
	Total	3657	488	0.13	1.00
2015	0.5	3832	2565	0.67	1.00
	2	3472	1399	0.40	1.00
	5	2774	912	0.33	1.00
	15	3187	654	0.21	1.00
	30	2566	287	0.11	1.00
	45	3364	174	0.05	1.00
	Total	19 195	5991	0.31	1.00

<sup>a</sup>Distance of mesotrione-susceptible *A. tuberculatus* female plants from the pollen-source block of mesotrione-resistant *A. tuberculatus*.

<sup>b</sup>Number of emerged and living plants ( $F_1$  individuals, resistant *A. tuberculatus* × susceptible *A. tuberculatus*) treated with 210 g ai ha<sup>-1</sup> (twice the mesotrione field rate) in intraspecific hybridization only.

<sup>c</sup>The frequency in  $F_1$  individuals was calculated from: frequency = number of emerged plants/number of living plants.


<sup>d</sup>Power analyses calculated for  $\alpha = 0.05$  using binomial probabilities.

While misdiagnosis of R *A. tuberculatus* and S *A. palmeri* hybrids with KASP assays may possibly be due to gene conversion, the chances of this happening at both loci is extremely low. Our results showed no evidence that allele conversion is happening at either locus, as the ALS-SNP and TBP-ITS (KASP assays) are in full agreement (Figure 1). The KASP assays greatly strengthen the case that heterozygous individuals are really hybrids and not artifacts of low levels of within-species

polymorphism. Further hybridization studies with KASP markers, especially the TBP-ITS assay, in other *Amaranthus* species are necessary for testing their general usefulness. We believe that the TBP-ITS marker has great potential for weed management decisions because the TBP-ITS marker robustly detects R *A. tuberculatus*, S *A. palmeri* and their hybrid.


In the field research site and adjacent areas, *A. palmeri* was not present prior to this experiment. Thus it is unlikely that pollen from male S *A. palmeri* could have outcrossed the female plants. Post-emergence mesotrione treatment reduced the number of putative hybrids by killing a majority of screened seedlings; however, mesotrione by itself was not a robust marker for detecting interspecific hybridization. Several hypotheses were raised to explain the high number of putative hybrids produced that were not mesotrione resistant and the high number of living putative hybrid after mesotrione treatment. First, mesotrione efficacy on *A. palmeri* varies with temperature (Godar *et al.*, 2015). In this interspecific study, mesotrione screening was performed under field conditions in June until August in 2015 and 2016, and the variable temperature might have influenced mesotrione efficacy. Second, multiple plants were grown in a single experimental unit. Thus false hybrids may have survived mesotrione treatment due to low herbicide coverage and therefore low herbicide dose. Third, the high number of non-hybrid seeds that were produced could be explained by facultative apomixis in female *A. palmeri* plants (Ribeiro *et al.*, 2014). The molecular mechanisms of apomictic seed formation in *A. palmeri* need further exploration.

It is possible that more hybrids were produced but the individuals were susceptible to mesotrione (at 1.75 times the mesotrione field rate). Therefore, the herbicide treatment could have masked the true number of hybrids in this study. Mesotrione treatment was performed to test the hypothesis of transfer of herbicide resistance between *Amaranthus* species. Nonetheless, all putative hybrid seedlings screened using molecular markers were either true hybrids (heterozygous for the S *A. palmeri* marker) or



**Figure 3.** Distance estimation of gene flow reduction frequency.

Distance (m) estimation of 50% and 90% frequency of gene flow ( $F_{GF}$ ) reduction in  $F_1$  individuals in eight directions of the intraspecific hybridization (mesotrione-resistant *Amaranthus tuberculatus* × mesotrione susceptible *A. tuberculatus*) field research experiment in 2014 (left) and 2015 (right).



**Figure 4.** Wind rose plots.

Wind rose plots demonstrating the average hourly (07:00 to 15:00) wind frequency (%) and wind speed ( $\text{m s}^{-1}$ ) grouped in  $22.5^\circ$  of direction (from which the wind is blowing) in a field experiment in 2014 (left) and 2015 (right). Wind data were recorded at 2 m above the soil surface and at the center of the pollen-source block of the field research experiment during the 6-week pollination period at the Haskell Agricultural Laboratory of the University of Nebraska–Lincoln. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)].

homozygous *S. A. palmeri*. No homozygous *R. A. tuberculatus* individuals were detected.

#### Intraspecific (*R. A. tuberculatus* × *S. A. tuberculatus*) hybridization

The transfer of herbicide resistance within populations of the same species is more likely to occur with high frequency (no genetic barrier) than between two different species (Sarangi *et al.*, 2017). Herbicide resistance traits are commonly used to measure hybridization as they can be easily screened for in  $F_1$  seedlings if they are autosomal dominant (Mallory-Smith *et al.*, 2015). Here, we accept our hypothesis as it was shown that under field research conditions mesotrione resistance alleles were transferred from *R. A. tuberculatus* to *S. A. tuberculatus* progeny through pollen. The frequency of gene flow was rapidly reduced within 45 m of the pollen source and varied with distance and direction, which was also documented with PMGF studies that used ALS (Liu *et al.*, 2012) and glyphosate resistance genes (Sosnoskie *et al.*, 2012; Ganie and Jhala, 2017; Sarangi *et al.*, 2017) in *Amaranthus* species. The mechanism of mesotrione resistance in *R. A. tuberculatus* is herbicide detoxification via cytochrome P450 alleles (Oliveira *et al.*, 2018c). Therefore, in this study it was not possible to use a molecular marker to detect mesotrione resistance in  $F_1$  seedlings due to the complexity of this herbicide resistance mechanism. Nonetheless, the modeling approach demonstrated that mesotrione resistance is inherited by the next generation under field research conditions. Oliveira *et al.* (2018a) documented that mesotrione resistance alleles in *R. A. tuberculatus* are polygenic, incompletely dominant and nuclear inherited, and this was also the case in other mesotrione-resistant *A. tuberculatus* populations from Illinois (Huffman *et al.*, 2015) and Iowa (Kohlhase *et al.*, 2018). The polygenic nature of mesotrione resistance traits suggests that resistance may evolve at a slower

pace within a population than single-gene dominant traits (Jasieniuk *et al.*, 1996; Huffman *et al.*, 2015).

The variability in the modeling approach demonstrated by the ME is probably related to the species biology, the phenotypic marker used and the complex polygenic nature of mesotrione resistance in this population. Variability within an experimental unit was expected as seeds were harvested from at least three *S. A. tuberculatus* plants in the receptor–donor block. Therefore, each  $F_1$  individual is likely to have a slightly different level of resistance, and it is possible that plants with either a high or a low resistance level were by chance sown in a determined experimental unit. It is also possible that ME of  $F_{GF}$  was higher in 2014 than 2015 because fewer replicates were used in 2015.

The mortality in  $F_1$  plants could be due to pollination of *S. A. tuberculatus* from adjacent areas as pollen can travel up to 800 m (Liu *et al.*, 2012), unintentional presence of *S. A. tuberculatus* in the pollen-donor block, an inability of heterozygous individuals to survive herbicide (twice the mesotrione field rate) or potential apomictic seed production by *S. A. tuberculatus*; this needs further exploration. Also, these limitations of the study and the complexity of mesotrione resistance could be the reason for there being no correlation between  $F_{GF}$  and wind frequency. We hypothesized that this result is probably due to the inconsistency of wind direction within a day and week and the variable flowering window of the species. Other studies of PMGF have shown that wind frequency or speed did not correlate with frequency of PMGF in *Echinochloa crus-galli* (Bagavathiannan and Norsworthy, 2014), *A. palmeri* (Sosnoskie *et al.*, 2009) and *Linum usitatissimum* (Jhala *et al.*, 2011). In general, PMGF studies are conducted with bare soil as a background, thus differences in  $F_{GF}$  and wind frequency found in this study versus others are expected. In this study, a soybean canopy (65 cm tall) was present, which creates different wind flow dynamics that could

have affected PMGF. When a soybean canopy is present, the wind flow would tend to be slowed down due to friction (aerodynamic resistance) with the canopy (Baldocchi *et al.*, 1983). Aerodynamic resistance to wind flow is a function of canopy height and increases with increasing canopy height. Therefore, pollen would not be able to travel as far for a given wind speed as over bare soil ground (Ganie and Jhala, 2017; Sarangi *et al.*, 2017), probably resulting in weaker correlation between  $F_{GF}$  and wind frequency.

## CONCLUSIONS

This research has demonstrated that PMGF may be a factor contributing to the evolution of herbicide resistance in *Amaranthus* species in the landscape. Even with low interspecific hybridization and rapid exponential decay in intraspecific hybridization with distance, PMGF carrying metabolism-based mesotrione resistance alleles occurs in weedy *Amaranthus* under field research conditions. This result is significant as *A. palmeri* and *A. tuberculatus* are prolific seed producers, obligate outcrossers, fast growing competitive, and NTSR can confer resistance to different herbicide SOA. Therefore, even at low frequencies, PMGF might have important evolutionary consequences in weedy *Amaranthus*.

## EXPERIMENTAL PROCEDURES

### Plant materials

The seeds from R and S *A. tuberculatus* populations were harvested from Platte County and Dixon County, NE, USA, respectively. The R *A. tuberculatus* has been previously characterized with 18-fold resistance to mesotrione compared with an S *A. tuberculatus* population (Oliveira *et al.*, 2017). The mechanism of herbicide resistance in the R *A. tuberculatus* population was described as incomplete dominance with multiple genes conferring enhanced mesotrione metabolism via cytochrome P450 enzymes (Kaundun *et al.*, 2017; Oliveira *et al.*, 2017, 2018a,b,c). The S *A. palmeri* population was collected from Lancaster County, NE, USA in 2001. In a preliminary study, this population was characterized as susceptible in the greenhouse with a 95% control value of 173 ( $\pm 62$ ) g ai ha<sup>-1</sup> of mesotrione. Seeds were soaked in water for a day before being manually sown in the field experiments.

### Field experiments

Field experiments were conducted in 2014 and 2015 in the same field at Haskell Agricultural Laboratory (42°23'1" N, 96°59'18" W) of the University of Nebraska–Lincoln at Concord, NE, USA. The soil type at the research site was loam with 2.7% organic matter and pH 7.6. The endemic weed species present at this site were *Setaria viridis*, *Chenopodium album*, *Abutilon theophrasti*, *A. tuberculatus* (mesotrione susceptible) and *A. retroflexus*. The soil was tilled, and glyphosate-resistant soybean was planted at 370 500 seeds ha<sup>-1</sup> in rows spaced 50 cm apart on 24 May 2014 and 25 May 2015.

The field experiment was conducted in an adapted concentric donor–receptor design, where the pollen-donor block was surrounded in eight directions by pollen-receptor blocks (Mallory-

Smith *et al.*, 2015). The pollen-donor block was a square (10 m × 10 m) and there were eight receptor blocks, each measuring 4 m wide and 45 m long (Figure 5). The total experimental area (pollen source and eight pollen-receptor blocks) was 1540 m<sup>2</sup>. Glufosinate (Liberty<sup>®</sup> 280 SL, BASF Agriculture, <https://agriculture.basf.com/>) was applied at 594 g ai ha<sup>-1</sup> to each section of the experimental area on soybean at cotyledon stage to provide weed- and crop-free areas prior to the onset of the experiment. Soybean was grown around the experimental area, which simulated the real field scenario and helped to suppress the endemic weeds in the experimental site (Figure 5). The 3-ha area external to the experimental site was sprayed with glyphosate (PowerMax<sup>®</sup>, Bayer Crop Science, <https://www.cropscience.bayer.com/>) at 1320 g ai ha<sup>-1</sup> for weed control, further reducing potential pollen contamination from the endemic *A. tuberculatus* and *A. retroflexus* population. Also, hoe weeding was performed within the experimental area for controlling undesired weed species that escaped herbicide treatment during the experiment period.

The *Amaranthus* species were manually sown on 2 June 2014 and 3 June 2015. This time was chosen because it is the natural starting germination window of *Amaranthus* species in north-eastern Nebraska. In the pollen-donor block, approximately 100 R *A. tuberculatus* seeds were transplanted 30 cm apart and 1 cm deep in the soil. A week later, the R *A. tuberculatus* population was sown again at 30 cm apart; therefore, R *A. tuberculatus* plants were spaced at a distance of 15 cm. Similarly, in the receptor



**Figure 5.** Aerial view of the field experiment.

Aerial view of the field experiment to quantify interspecific and intraspecific pollen-mediated gene flow from mesotrione-resistant *Amaranthus tuberculatus* (R) to mesotrione-susceptible *Amaranthus palmeri* and *A. tuberculatus* (S). The R *A. tuberculatus* was planted in the center of the field in a 10 m × 10 m pollen-source block. The pollen-receptor blocks were divided into eight cardinal direction blocks (N, NE, E, SE, S, SW, W, NW, 4 m × 45 m each). The S *A. palmeri* and S *A. tuberculatus* were planted at each pollen-receptor block at 0.5, 2, 5, 15, 30 and 45 m from the pollen-source block. At each distance in the pollen-receptor block, *A. palmeri* and *A. tuberculatus* were planted in 3 and 1 linear meters, respectively. [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)].



blocks, 100 seeds of *S. A. tuberculatus* and *S. A. palmeri* were also sown in at an interval of a week spaced 15 cm apart. The receptor block plants were sown at six distances (0.5, 2, 5, 15, 30 and 45 m) in eight different blocks, including the cardinal (north, south, east, west) and the ordinal (north-east, north-west, south-east, south-west) directions (Figure 5). At each distance and direction, *S. A. tuberculatus* and *S. A. palmeri* were sown 1 and 3 m wide (four linear meters), respectively. Inside the pollen-donor area, *S. A. palmeri* was also sown in two linear meters to simulate a 'worst-case' scenario, where *S. A. palmeri* was surrounded by a dense population of *R. A. tuberculatus* plants. The week's interval between planting was allowed to increase synchrony of the flowering period and therefore the chance of hybridization between species.

After the *A. tuberculatus* and *A. palmeri* seedlings had germinated, thinning was performed to reduce plant competition, but dense populations of *A. tuberculatus* and *A. palmeri* were maintained in the study (approximately 30 plants per linear meter). Male *S. A. tuberculatus* and *S. A. palmeri* plants were visually detected on a daily-basis prior to pollen shedding and removed from the pollen-receptor and pollen-donor blocks (*S. A. tuberculatus*). Also, in the pollen-donor block, female *R. A. tuberculatus* plants were similarly screened and removed. This procedure was performed to maximize outcrossing between male *R. A. tuberculatus* and female *S. A. tuberculatus* and *S. A. palmeri* plants (Figure S4). Otherwise, pollen competition may have occurred, reducing the chances for distant pollen dispersal in the study (Liu *et al.*, 2012).

Both *Amaranthus* species in these studies displayed long flowering periods from mid July to late August in both 2014 and 2015. The height of *Amaranthus* species at flowering varied from 1 to 2 m. A weather station (61622 Vantage Pro2 Plus, Davis Instruments, <https://www.davisinstruments.com/>) was placed at 2 m above the soil surface in the center of this experiment. Hourly meteorological data on temperature, humidity and precipitation were recorded (Table S1). The wind speed and direction data were recorded from 07:00 to 15:00 due to the likelihood of *Amaranthus* species shedding pollen during that period (Sosnoskie *et al.*, 2012).

At maturity (late September), inflorescences of ten *S. A. palmeri* and three *S. A. tuberculatus* plants were harvested from each distance and direction. Harvesting was performed moving inward, starting at 45 m and progressing to 0.5 m for each species to reduce seed contamination. Mature inflorescences of species at each direction and distance were cleaned, labeled and bagged individually, then stored at 4°C to disrupt potential seed dormancy until they could be used in the herbicide resistance screening. The harvested seeds were termed putative hybrids (*R. A. tuberculatus* × *S. A. palmeri*) and F<sub>1</sub> (*R. A. tuberculatus* × *S. A. tuberculatus*) individuals to describe interspecific and intraspecific hybridization, respectively.

#### Interspecific hybridization (*A. tuberculatus* × *A. palmeri*) species diagnostic marker

Three markers (one phenotypic and two genotypic) were combined to detect hybrids. First, mesotrione resistance was used as a phenotypic marker to increase the probability of finding hybrids. Second, plants that survived mesotrione treatment were genotyped using two KASP assays to determine whether each individual was a true hybrid between *S. A. palmeri* and *R. A. tuberculatus* (Figure S5).

**Mesotrione resistance marker.** Hybrid seeds collected from the field study were planted separately in plastic trays

(51 cm × 38 cm × 10 cm) containing potting mix (Miracle-Gro®, ScottsMiracle-Gro, <https://scottsmiracleagro.com/>) and evaluated for mesotrione resistance.

The mesotrione screening on putative hybrids was performed outdoors in the summer period of 2015 and 2016 at the Haskell Agricultural Laboratory. There was a high variability in hybrid germination in trays. For example, in each tray, putative hybrids varied from 0 to 1296 seedlings with an average of 209 and 63 seedlings per tray in 2014 and 2015, respectively. There were eight to nine replications of each experimental unit (plastic trays) in 2014 and 11 replications from the 2015 study. A total of 104 492 putative hybrids seedlings were screened in all distances and directions for the study in 2014 and 2015. Putative hybrids were sprayed at 5–8 cm tall with 1.75 times (interspecific hybridization only) the mesotrione field rate (105 g ai ha<sup>-1</sup>; Callisto®, Syngenta Crop Protection, <http://www.syngentacropprotection.com/>). Mesotrione was applied as described in Oliveira *et al.* (2017). Hybrids were assessed as dead or alive 21 days after mesotrione application, and the number of surviving plants was recorded. In general, surviving hybrid seedlings showed 40% to 85% injury to mesotrione, using a scale rate of 0% to 100% (no injury to plant death), as described by Oliveira *et al.* (2018a,b). Leaf tissue of surviving hybrid seedlings was collected and stored at –80°C to be used for KASP assay analysis.

**Competitive allele-specific PCR (KASP) assays.** DNA was extracted from four parental *S. A. palmeri* seedlings, four parental *R. A. tuberculatus* plants, and 86 putative hybrids that survived mesotrione treatment at 1.75 times mesotrione field rate (phenotypic marker). DNA was extracted using a modified cetyl trimethylammonium bromide method (Doyle, 1987) described in detail by Patterson *et al.* (2017). All DNA was then diluted to 5 ng μl<sup>-1</sup> for the KASP assays.

Two KASP assays were used to determine whether individuals had *S. A. palmeri*, *R. A. tuberculatus* or hybrid (heterozygous) genotypes. The first SNP was located at base pair 678 in the ALS coding sequence and has been used previously as a RFLP marker (Tranel *et al.*, 2002) and as a KASP marker (Küpper *et al.*, 2017). The second marker is a 2-bp polymorphism in the ITS of the ribosomal coding region (TBP-ITS), designed to distinguish *A. palmeri* from eight other *Amaranthus* species (Table S4). This polymorphism is at base pairs 496 and 497 in the ITS sequence from *A. palmeri* accession KP318856.1 of the National Center for Biotechnology Information (NCBI) nucleotide database (Figure 6). At this locus, *A. palmeri* has an adenosine (A) followed by a guanine (G), while all other species in the *Amaranthus* genus have a cytosine (C) followed by an adenosine (A).

The protocol detailed in Küpper *et al.* (2017) was used for the ALS marker. In brief, the ALS-SNP assay used the following primers: *A. tuberculatus* forward primer, 5'-GAAGGTGACCAAGTT-CATGCTAAAAAGAAAGCTTCCTTAACAATTCTAGGG-3' (FAM Tag underlined); *A. palmeri* forward primer, 5'-GAAGTTCGGAGT-CAACGGATTAAAAAGAAAGCTTCCTTAACAATTCTAGGA-3' (HEX Tag underlined); universal reverse primer, 5'-GTTGAGGTAAGTC-GATCCACTACTA AGC-3'. For the TBP-ITS, two species-diagnostic forward primers were developed that were identical, except for the final two 3' nucleotides, which pair with base pairs 496 and 497. Additionally, each forward primer was tagged at its 5' end with nucleotides that are specific for either a HEX- or FAM-labeled oligo that comes pre-mixed in KASP Master Mix (LGC Genomics, <https://www.lgcgroup.com/>). For the TBP-ITS assay we used the following primers: *A. tuberculatus* forward primer, 5'-GAAGTCG-GAGTCAACGGATTTCGGCGTGGATGGCCTAAAACA-3' (FAM Tag



non-linear models (package *gnm*) in R statistical software (Turner and Firth, 2015). Fourteen models were tested for describing the intraspecific frequency of gene flow (Table S2). The Akaike information criterion (AIC) was used to select the top model to describe the frequency of intraspecific gene flow in *A. tuberculatus*. According to the AIC criterion, the top model has the lowest AIC value (Oliveira *et al.*, 2018a,b).

Based on the AIC, the top model was a double exponential decay (Tables S2 and S3), with the  $F_{GF}$  in  $F_1$  varying with distance from the pollen-source block, the direction of the pollen-receptor block and the year of the experiment:

$$\text{logit}(p_i) = l_0 + \exp[l_1 + d_1 \times \text{Distance}] + \exp[l_2(\text{Direction:Year}) + d_2(\text{Direction:Year}) \times \text{Distance}] \quad (4)$$

where  $p_i$  is the  $F_{GF}$  in the  $i$ th observation,  $l_0$  is the overall intercept,  $l_1$  and  $l_2$  are intercepts of the first and second exponential instances and  $d_1$  and  $d_2$  are the decay rates. In this model,  $l_2$  and  $d_2$  vary with direction and year.

The distances for 50% and 90%  $F_{GF}$  reduction were estimated from equation (4) in each of the eight directions of the pollen-receptor block. The ME was calculated to test the goodness of fit of the top model for each direction (Werle *et al.*, 2014):

$$\text{ME} = 1 - \frac{\sum_{i=1}^n (O_i - P_i)^2}{\sum_{i=1}^n (O_i - \bar{O}_i)^2} \quad (5)$$

where  $n$  is the number of data points,  $O_i$  is the observed value,  $P_i$  is the predicted value and  $\bar{O}_i$  is the mean observed value. The ME values range from  $-\infty$  to 1, with values closer to 1 indicating better predictions.

## Power analysis

A power analysis for binomial probabilities was performed to determine the statistical precision of hybridization with the sample size used in this experiment. The theoretical values used were 0.5% and 1% at  $\alpha = 0.05$  for interspecific and intraspecific hybridization, respectively. The theoretical hybridization frequencies were compared with the observed hybridization frequencies as described in Jhala *et al.* (2011).

## ACKNOWLEDGEMENTS

The authors would like to thank CAPES (Brazilian Government Foundation) Proc. no. 9112-13-8 for financial support to the first author of this paper. In addition, we thank Dr Jon Scott, Sérgio Oliveira, Ícaro Freitas, Amanda Winstead, Kyle Kardell, Nick Luhr, Tarin Quin and Brad Steffen for their assistance with the field experiments.

## CONFLICT OF INTEREST

The authors declare no conflict of interest.

## SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article.

**Figure S1.** Frequency of gene flow predicted with a double exponential model.

**Figure S2.** Wind rose of wind frequency (%) and wind speed ( $\text{m s}^{-1}$ ) in 2014.

**Figure S3.** Wind rose of wind frequency (%) and wind speed ( $\text{m s}^{-1}$ ) in 2015.

**Figure S4.** Inflorescences of *Amaranthus tuberculatus* (male and female) and *Amaranthus palmeri* (female).

**Figure S5.** Methods for detecting the interspecific hybrids.

**Table S1.** Weather data (30-year average) at the experimental site.

**Table S2.** Candidate models for describing intraspecific hybrids.

**Table S3.** Coefficient estimation from a double-exponential decay model.

**Table S4.** *Amaranthus* species accession from the alignment of the internal transcribed spacer region.

## REFERENCES

- Bagavathiannan, M.V. and Norsworthy, J.K. (2014) Pollen-mediated transfer of herbicide resistance in *Echinochloa crus-galli*. *Pest Manag. Sci.* **70**, 1425–1431.
- Baldocchi, D.D., Verma, S.B. and Rosenberg, N.J. (1983) Characteristics of air flow above and within soybean canopies. *Boundary-Layer Meteorol.* **25**, 43–54.
- Bernards, M.L., Crespo, R.J., Kruger, G.R., Gaussoin, R. and Tranel, P.J. (2012) A waterhemp (*Amaranthus tuberculatus*) population resistant to 2,4-D. *Weed Sci.* **60**, 379–384.
- Culpepper, A.S. (2006) Glyphosate-induced weed shifts. *Weed Technol.* **20**, 277–281.
- Délye, C. (2013a) Unravelling the genetic bases of non-target-site-based resistance (NTSR) to herbicides: a major challenge for weed science in the forthcoming decade. *Pest Manag. Sci.* **69**, 176–187.
- Délye, C., Jasieniuk, M., Rie, V. and Corre, L. (2013b) Deciphering the evolution of herbicide resistance in weeds. *Trends Genet.* **29**, 649–658.
- Doyle, J.J. (1987) A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochem. Bull.* **19**, 11–15.
- Duhoux, A., Carrère, S., Gouzy, J., Bonin, L. and Délye, C. (2015) RNA-seq analysis of rye-grass transcriptomic response to an herbicide inhibiting acetolactate-synthase identifies transcripts linked to non-target-site-based resistance. *Plant Mol. Biol.* **87**, 473–487.
- Ellstrand, N.C., Whitkus, R., Rieseberg, L.H. and Anderson, W.W. (1996) Distribution of spontaneous plant hybrids. *Proc. Natl Acad. Sci. USA* **93**, 5090–5093.
- Evans, A.F., O'Brien, S.R., Ma, R., Hager, A.G., Riggins, C.W., Lambert, K.N. and Riechers, D.E. (2017) Biochemical characterization of metabolism-based atrazine resistance in *Amaranthus tuberculatus* and identification of an expressed GST associated with resistance. *Plant Biotechnol. J.* **15**, 1238–1249.
- Figueiredo, M.R., Leibhart, L.J., Reicher, Z.J., Tranel, P.J., Nissen, S.J., Westra, P., Bernards, M.L., Kruger, G.R., Gaines, T.A. and Jugulam, M. (2017) Metabolism of 2,4-dichlorophenoxyacetic acid contributes to resistance in a common waterhemp (*Amaranthus tuberculatus*) population. *Pest Manag. Sci.* **74**, 2356–2362.
- Franssen, A.S., Skinner, D.Z., Al-Khatib, K., Horak, M.J. and Kulakow, P.A. (2001a) Interspecific hybridization and gene flow of ALS resistance in *Amaranthus* species. *Weed Sci.* **49**, 598–606.
- Franssen, A.S., Skinner, D.Z., Al-Khatib, K. and Horak, M.J. (2001b) Pollen morphological differences in *Amaranthus* species and interspecific hybrids. *Weed Sci.* **49**, 732–737.
- Gaines, T.A., Ward, S.M., Bukun, B., Preston, C., Leach, J.E. and Westra, P. (2012) Interspecific hybridization transfers a previously unknown glyphosate resistance mechanism in *Amaranthus* species. *Evol. Appl.* **5**, 29–38.
- Gaines, T.A., Lorentz, L., Figge, A. *et al.* (2014) RNA-Seq transcriptome analysis to identify genes involved in metabolism-based diclofop resistance in *Lolium rigidum*. *Plant J.* **78**, 865–876.
- Ganie, Z.A. and Jhala, A.J. (2017) Modelling pollen-mediated gene flow from glyphosate-resistant to -susceptible giant ragweed (*Ambrosia trifida* L.) under field conditions. *Sci. Rep.* **7**, 17067.
- Godar, A.S., Varanasi, V.K., Nakka, S., Prasad, P.V.V., Thompson, C.R. and Mithila, J. (2015) Physiological and molecular mechanisms of differential sensitivity of Palmer amaranth (*Amaranthus palmeri*) to mesotrione at varying growth temperatures. *PLoS ONE* **10**, e0126731.
- Grant, W.F. (1959) Cytogenetics studies in *Amaranthus*: II. Natural interspecific hybridization between *Amaranthus dubius* and *A. spinosus*. *Can. J. Bot.* **37**, 1063–1070.
- Greizerstein, E.J. and Poggio, L. (1995) Meiotic studies of spontaneous hybrids of *Amaranthus*: genome analysis. *Plant Breed.* **114**, 448–450.
- Hartigan, J.A. and Wong, M.A. (1979) Algorithm AS 136: a K-means clustering algorithm. *J. R. Stat. Soc. Ser. C Appl. Stat.* **28**, 100–108.

- Hartzler, R.G., Battles, B.A. and Nordby, D. (2004) Effect of common waterhemp (*Amaranthus rudis*) emergence date on growth and fecundity in soybean. *Weed Sci.* **52**, 242–245.
- Hausman, N.E., Singh, S., Tranel, P.J., Riechers, D.E., Kaundun, S.S., Polge, N.D., Thomas, D.A. and Hager, A.G. (2011) Resistance to HPPD-inhibiting herbicides in a population of waterhemp (*Amaranthus tuberculatus*) from Illinois, USA. *Pest Manag. Sci.* **67**, 258–261.
- He, C., Holme, J. and Anthony, J. (2014) SNP genotyping: the KASP assay. In *Crop Breeding. Methods in Molecular Biology (Methods and Protocols)* (Fleury, D. and Whitford, R., eds). New York: Humana Press, pp. 75–86.
- Huffman, J., Hausman, N.E., Hager, A.G., Riechers, D.E. and Tranel, P.J. (2015) Genetics and inheritance of nontarget-site resistances to atrazine and mesotrione in a waterhemp (*Amaranthus tuberculatus*) population from Illinois. *Weed Sci.* **63**, 799–809.
- Jasieniuk, M., Brulé-Babel, A.L. and Morrison, I.N. (1996) The evolution and genetics of herbicide resistance in weeds. *Weed Sci.* **44**, 176–193.
- Jhala, A.J., Bhatt, H., Topinka, K. and Hall, L.M. (2011) Pollen-mediated gene flow in flax (*Linum usitatissimum* L.): can genetically engineered and organic flax coexist? *Heredity*, **106**, 557–566.
- Kaundun, S.S., Hutchings, S.-J., Dale, R.P., Howell, A., Morris, J.A., Kramer, V.C., Shivrain, V.K. and McIndoe, E. (2017) Mechanism of resistance to mesotrione in an *Amaranthus tuberculatus* population from Nebraska, USA. *PLoS ONE*, **12**, e0180095.
- Kohlhase, D., Edwards, J. and Owen, M. (2018) Inheritance of 4-hydroxyphenylpyruvate dioxygenase inhibitor herbicide resistance in *Amaranthus tuberculatus*. *Plant Sci.* **274**, 360–368. <https://doi.org/10.1016/j.plantsci.2018.06.004>.
- Kohrt, J.R., Sprague, C.L., Swathi Nadakuduti, S. and Douches, D. (2017) Confirmation of a three-way (Glyphosate, ALS, and Atrazine) herbicide-resistant population of Palmer amaranth (*Amaranthus palmeri*) in Michigan. *Weed Sci.* **65**, 327–338.
- Küpper, A., Borgato, E.A., Patterson, E.L., Netto, A.G., Nicolai, M., Carvalho, S.J.P., De, Nissen, S.J., Gaines, T.A. and Christoffoleti, P.J. (2017) Multiple resistance to glyphosate and acetolactate synthase inhibitors in Palmer amaranth (*Amaranthus palmeri*) identified in Brazil. *Weed Sci.* **65**, 317–326.
- Küpper, A., Peter, F., Zöllner, P., Lorentz, L., Tranel, P.J., Beffa, R. and Gaines, T.A. (2018) Tembotrione detoxification in 4-hydroxyphenylpyruvate dioxygenase (HPPD) inhibitor-resistant Palmer amaranth (*Amaranthus palmeri* S. Wats.). *Pest Manag. Sci.* **15**, 1238–1249.
- Liu, J., Davis, A.S. and Tranel, P.J. (2012) Pollen biology and dispersal dynamics in waterhemp (*Amaranthus tuberculatus*). *Weed Sci.* **60**, 416–422.
- Liu, C., Bridges, M.E., Kaundun, S.S., Glasgow, L., Owen, M.D. and Neve, P. (2017) A generalised individual-based algorithm for modelling the evolution of quantitative herbicide resistance in arable weed populations. *Pest Manag. Sci.* **73**, 462–474.
- Mallory-Smith, C., Hall, L.M. and Burgos, N.R. (2015) Experimental methods to study gene flow. *Weed Sci.* **63**, 12–22.
- Murphy, B.P., Plewa, D.E., Phillippi, E., Bissonnette, S.M. and Tranel, P.J. (2017) A quantitative assay for *Amaranthus palmeri* identification. *Pest Manag. Sci.* **73**, 2221–2224. <https://doi.org/10.1002/ps.4632>.
- Nakka, S., Godar, A.S., Wani, P.S., Thompson, C.R., Peterson, D.E., Roelofs, J. and Jugulam, M. (2017) Physiological and molecular characterization of hydroxyphenylpyruvate dioxygenase (HPPD)-inhibitor resistance in Palmer amaranth (*Amaranthus palmeri* S.Wats.). *Front. Plant Sci.* **8**, 555.
- Nandula, V.K., Wright, A.A., Bond, J.A., Ray, J.D., Eubank, T.W. and Molin, W.T. (2014) EPSPS amplification in glyphosate-resistant spiny amaranth (*Amaranthus spinosus*): a case of gene transfer via interspecific hybridization from glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*). *Pest Manag. Sci.* **70**, 1902–1909.
- Oliveira, M.C., Jhala, A.J., Gaines, T., Irmak, S., Amundsen, K., Scott, J.E. and Knezevic, S.Z. (2017) Confirmation and control of HPPD-inhibiting herbicide-resistant waterhemp (*Amaranthus tuberculatus*) in Nebraska. *Weed Technol.* **31**, 67–79.
- Oliveira, M.C., Gaines, T.A., Dayan, F.E., Patterson, E.L., Jhala, A.J. and Knezevic, S.Z. (2018c) Reversing resistance to tembotrione in an *Amaranthus tuberculatus* (var. *rudis*) population from Nebraska, USA with cytochrome P450 inhibitors. *Pest Manag. Sci.* **74**, 2296–2305. <https://doi.org/10.1002/ps.4697>.
- Oliveira, M.C., Gaines, T.A., Jhala, A.J. and Knezevic, S.Z. (2018a) Inheritance of mesotrione resistance in an *Amaranthus tuberculatus* (var. *rudis*) population from Nebraska, USA. *Front. Plant Sci.* **9**, 1–12. <https://doi.org/10.3389/fpls.2018.00060>.
- Oliveira, M.C., Pereira, G.A.M., Ferreira, E.A., Santos, J.B., Knezevic, S.Z. and Werle, R. (2018b) Additive design: the concept and data analysis. *Weed Res.* **58**, 338–347. <https://doi.org/10.1111/wer.12317>.
- Owen, M.D. (2008) Weed species shifts in glyphosate-resistant crops. *Pest Manag. Sci.* **64**, 377–387.
- Pan, L., Gao, H., Xia, W., Zhang, T. and Dong, L. (2016) Establishing a herbicide-metabolizing enzyme library in *Beckmannia syzigachne* to identify genes associated with metabolic resistance. *J. Exp. Bot.* **67**, 1745–1757.
- Patterson, E.L., Fleming, M.B., Kessler, K.C., Nissen, S.J. and Gaines, T.A. (2017) A KASP genotyping method to identify northern watermilfoil, Eurasian watermilfoil, and their interspecific hybrids. *Front. Plant Sci.* **8**, 752.
- Refsell, D.E. and Hartzler, R.G. (2009) Effect of tillage on common waterhemp (*Amaranthus rudis*) emergence and vertical distribution of seed in the soil. *Weed Technol.* **23**, 129–133.
- Ribeiro, D.N., Pan, Z., Duke, S.O., Nandula, V.K., Baldwin, B.S., Shaw, D.R. and Dayan, F.E. (2014) Involvement of facultative apomixis in inheritance of EPSPS gene amplification in glyphosate-resistant *Amaranthus palmeri*. *Planta*, **239**, 199–212.
- Sarangi, D., Tyre, A.J., Patterson, E.L., Gaines, T.A., Irmak, S., Knezevic, S.Z., Lindquist, J.L. and Jhala, A.J. (2017) Pollen-mediated gene flow from glyphosate-resistant common waterhemp (*Amaranthus rudis* Sauer): consequences for the dispersal of resistance genes. *Sci. Rep.* **7**, 44913.
- Sauer, J. (1957) Recent migration and evolution of the dioecious amaranths. *Evolution*, **11**, 11–31.
- Sauer, J.D. (1967) The grain amaranths and their relatives: a revised taxonomic and geographic survey. *Ann. Missouri Bot. Gard.* **54**, 103.
- Sosnoskie, L.M., Webster, T.M., Dales, D., Rains, G.C., Grey, T.L. and Culpepper, A.S. (2009) Pollen grain size, density, and settling velocity for Palmer amaranth (*Amaranthus palmeri*). *Weed Sci.* **4**, 404–409.
- Sosnoskie, L.M., Webster, T.M., Kichler, J.M., Macrae, A.W., Grey, T.L. and Culpepper, A.S. (2012) Pollen-mediated dispersal of glyphosate-resistance in Palmer amaranth under field conditions. *Weed Sci.* **60**, 366–373.
- Steckel, L.E. (2007) The dioecious *Amaranthus* spp.: here to stay. *Weed Technol.* **21**, 567–570.
- Steinau, A.N., Skinner, D.Z. and Steinau, M. (2003) Mechanism of extreme genetic recombination in weedy *Amaranthus* hybrids. *Weed Sci.* **51**, 696–701.
- Tranel, P.J., Wassom, J.J., Jeschke, M.R. and Rayburn, A.L. (2002) Transmission of herbicide resistance from a monoecious to a dioecious weedy *Amaranthus* species. *Theor. Appl. Genet.* **105**, 674–679.
- Tranel, P.J., Riggins, C.W., Bell, M.S. and Hager, A.G. (2011) Herbicide resistances in *Amaranthus tuberculatus*: a call for new options. *J. Agric. Food Chem.* **59**, 5808–5812.
- Trucco, F., Jeschke, M.R., Rayburn, A.L. and Tranel, P.J. (2005) Promiscuity in weedy amaranths: high frequency of female tall waterhemp (*Amaranthus tuberculatus*) × smooth pigweed (*A. hybridus*) hybridization under field conditions. *Weed Sci.* **53**, 46–54.
- Trucco, F., Tatum, T., Robertson, K.R., Lane Rayburn, A. and Tranel, P.J. (2006) Characterization of waterhemp (*Amaranthus tuberculatus*) × Smooth Pigweed (*A. hybridus*) F1 hybrids. *Weed Technol.* **20**, 14–22.
- Trucco, F., Zheng, D., Woodyard, A.J., Walter, J.R., Tatum, T.C., Lane Rayburn, A. and Tranel, P.J. (2007) Nonhybrid progeny from crosses of dioecious amaranths: implications for gene-flow research. *Weed Sci.* **55**, 119–122.
- Trucco, F., Tatum, T., Rayburn, K.R., Lane Rayburn, A. and Tranel, P.J. (2009) Out of the swamp: unidirectional hybridization with weedy species may explain the prevalence of *Amaranthus tuberculatus* as a weed. *New Phytol.* **184**, 819–827.
- Turner, H. and Firth, D. (2015) Generalized nonlinear models in R: an overview of the gnm package generalized linear models preamble. (*R package version 1.0-8*) (<http://CRAN.R-project.org/package=gnm>).
- Ward, S.M., Webster, T.M. and Steckel, L.E. (2013) Palmer amaranth (*Amaranthus palmeri*): a review. *Weed Technol.* **27**, 12–27.
- Wassom, J.J. and Tranel, P.J. (2005) Amplified fragment length polymorphism-based genetic relationships among weedy *Amaranthus* species. *J. Hered.* **96**, 410–416.

- Webster, T.M. and Nichols, R.L.** (2012) Changes in the prevalence of weed species in the major agronomic crops of the southern United States: 1994/1995 to 2008/2009. *Weed Sci.* **60**, 145–157.
- Werle, R., Schmidt, J.J., Laborde, J., Tran, A., Creech, C.F. and Lindquist, J.L.** (2014) Shattercane X ALS-tolerant sorghum F1 hybrid and shattercane interference in ALS-tolerant sorghum. *J. Agric. Sci.* **6**, 159–165.
- Wetzel, D.K., Horak, M.J. and Skinner, D.Z.** (1999a) Use of PCR-based molecular markers to identify weedy *Amaranthus* species. *Weed Sci.* **47**, 518–523.
- Wetzel, D.K., Horak, M.J., Skinner, D.Z. and Kulakow, P.A.** (1999b) Transfer of herbicide resistance traits from *Amaranthus palmeri* to *Amaranthus rudis*. *Weed Sci.* **47**, 538–543.
- Wright, A.A., Molin, W.T. and Nandula, V.K.** (2016) Distinguishing between weedy *Amaranthus* species based on intron 1 sequences from the 5-enolpyruvylshikimate-3-phosphate synthase gene. *Pest Manag. Sci.* **72**, 2347–2354.