

## **Impact of annual precipitation on corn yield in a long-term tillage and N rate study**


**Javed Iqbal, Charles Shapiro, Ankit Shekhar**  
University of Nebraska-Lincoln

## Goals of this presentation

- Know the impact of annual precipitation on corn yield under certain management practices
- Decide if your farm management practices can be adjusted with changing precipitation patterns

# Crop Production Clinics

**N** EXTENSION



# Crop Production Clinics

**N** EXTENSION

2011 Flood




# Crop Production Clinics

## 2012 Drought

### U.S. Drought Monitor: NEBRASKA

December 4, 2012



#### Drought Severity



Week	Nothing	D0-D4	D1-D4	D2-D4	D3-D4	D4
Nov 27, 2012	0.00	100.00	100.00	100.00	96.15	77.46
Dec 4, 2012	0.00	100.00	100.00	100.00	96.15	77.46

# Crop Production Clinics

**N** EXTENSION

## 2019 Flood




# Crop Production Clinics



## 2020 Drought *U.S. Drought Monitor* Nebraska

December 22, 2020  
(Released Thursday, Dec. 24, 2020)  
Valid 7 a.m. EST



***Intensity:***

- None
- D0 Abnormally Dry
- D1 Moderate Drought
- D2 Severe Drought
- D3 Extreme Drought
- D4 Exceptional Drought

*The Drought Monitor focuses on broad-scale conditions. Local conditions may vary. For more information on the Drought Monitor, go to <https://droughtmonitor.unl.edu/About.aspx>*

***Author:***

Adam Hartman  
NOAA/NWS/NCEP/CPC



[droughtmonitor.unl.edu](https://droughtmonitor.unl.edu)

## Objectives

Use Standardized Precipitation Evapotranspiration Index (SPEI) to

- Determine if SPEI helps differentiate the crop yield with tillage, nitrogen rate, and crop rotation across the index range, and
- Find out if SPEI helps in making management decisions with precipitation regimes



# Crop Production Clinics



## Methods

- Study site: Haskell Ag Lab, Northeast NE
- Study years: 1986-now (34 years)
- Study design: split-split plot
- Tillage treatments: No-Till (NT), Spring/fall plow (PL), Disk (DK)
- Crop rotation: Corn-Soybean (CS), Continuous Corn (CC)
- Nitrogen rates: 0, 40, 80, 120, and 160 kg N ha<sup>-1</sup>
- Nitrogen source: Ammonium Nitrate




# Crop Production Clinics

Standardized Precipitation Evapotranspiration Index (SPEI) and number of years in each category for 31 years


	Climatic Moisture categories	SPEI-5Sep OR SPEI-12Sep	Number of years	
Wet	Extremely wet	$\geq 2.00$	0	
	Severely wet	1.50–1.99	2 (1993, 2014)	(6 of 31 times, 19% probability)
	Moderately wet	1.00-1.49	4 (1992, 1996, 2010, 2015)	
Normal	Normal	0.99 to $-0.99$	19	(19 of 31 times, 61% probability)
Dry	Moderate drought	$-1.00$ to $-1.49$	3 (1990, 2000, 2002)	
	Severe drought	$-1.50$ to $-1.99$	1 (1989)	(6 of 31 times, 19% probability)
	Extreme drought	$\leq -2.00$	2 (1988, 2012)	

## Corn yield response to N rate



Shekhar and Shapiro et al. 2019

## Corn yield response to tillage



- Dry years
- Normal years
- Wet years
- Overall years

UNIVERSITY OF NEBRASKA-LINCOLN

**CROPWATCH**

Home Weather (GDD & ET) Info & Resources Crops Management Related Topics Archives

### Strategic Tillage for the Improvement of No-Till Cropping Systems

JANUARY 21, 2020

Charles Wortmann - Extension Soil and Nutrient Management Specialist | Humberto Blanco - Professor of Soil Science


Occasional tillage (OT) of long-term no-till (also called one-time or strategic tillage) might be desired once in more than 5 or 10 years according to need for integrated weed management, fracturing a compaction layer, incorporating a soil amendment such as lime or manure, reducing vertical stratification of nutrient availability, increasing soil organic matter to greater depth, or reducing crop residue accumulation. The type of tillage for OT should be specific to the objective of the OT. Some of the early OT research was done in Nebraska including a 5-year study at the High Plains Agricultural Laboratory (HPAL) near Sidney using moldboard plow tillage and three 5-year trials in eastern Nebraska in which five OT practices were compared. However, there has been much additional study elsewhere during the past decade. Dozens of other multi-year trials have been conducted including trials in Australia, Brazil, Canada, Spain, and Turkey, as well as in Indiana, Kansas, Kentucky, Oregon, Texas, and Wyoming.

#### Negative, neutral and positive effects

An early concern with OT of no till was that benefits to soil properties and productivity gained from continuous no-till would be lost with a single or infrequent tillage practice. The findings from the above studies consistently show that such negative effects are highly unlikely, generally of less than 1-year duration and of little agronomic significance. The two consistent negative effects of OT are the cost of the tillage and increased risk of erosion until crop residue or residue cover of the soil is established.

Shekhar and Shapiro et al. 2019

## Corn yield response to crop rotation



## Tillage, rotation and N rate recommendations based on SPEI at rainfed silt loam soil

Tillage, rotation, and N rate recommendations based on a meteorological drought index (SPEI) analysis over 31 years at a rainfed silt loam site in Nebraska, US if precipitation index could be predicted.

Recommendations			
Moisture Conditions	Tillage	Rotation	N rate
Extremely/Severely Dry ( $\text{SPEI} \leq -1.5$ )	No-Till	Continuous Corn (C-C)	-
Moderately Dry ( $-1 \geq \text{SPEI} \geq -1.49$ )	No-Till	Corn-Soybean (C-S) rotation	Maximum of $120 \text{ kg ha}^{-1}$
Normal ( $0.99 \geq \text{SPEI} \geq -0.99$ )	No-Till	Corn-Soybean (C-S) rotation	$120 \text{ kg ha}^{-1}$
Wet ( $\text{SPEI} \geq 1$ )	Plow or Disk	Both C-C and C-S	$150 \text{ kg ha}^{-1}$ for C-S and $220 \text{ kg ha}^{-1}$ for C-C

## Take Home Message

- **N Rate:** Under normal and dry conditions, (25 of 31 times, 81% probability) optimum NR of 120 kg N ha<sup>-1</sup> can be more profitable for a rainfed silt-loam soil.
- **Tillage:** Under normal and dry conditions (25 of 31 times, 81% probability), No-till enhances crop yield, profits, provides soils benefits, and can help to protect water quality.
- **Crop Rotation:** Rotation (C–S) under normal and wet conditions (25 of 31 times, 81% probability) increase yields.


# Crop Production Clinics

**N** EXTENSION



Javed Iqbal

 [jiqbal2@unl.edu](mailto:jiqbal2@unl.edu)

 402-472-1432

 **THANKS**